

PUBLISHED BY THE OIKOS EDITORIAL OFFICE

# NORDIC JOURNAL OF BOTANY

*Gymnosporia thyrsoflora* comb. nov.  
(Celastraceae), a correct name to replace  
*G. graciliramula* from southwest China

Wen-Bin Yu, Hong Wang and De-Zhu Li

DOI: 10.1111/j.1756-1051.2013.01672.x,  
Volume 31, Issue 6, Pages 746–747


If you wish to order reprints of this article,  
please follow the instructions [here](#)

## EMAIL ALERTS

Receive free email alerts and stay up-to-date on what is published  
in Nordic Journal of Botany – [click here](#)

## CONNECT WITH US

Follow Nordic Journal of Botany on [Twitter](#) and [Facebook](#)



Submit your next paper to NJBOT online at <http://mc.manuscriptcentral.com/njbot>

Subscribe to Nordic Journal of Botany and stay up-to-date with a wide variety of  
new findings of relevance to the botanical biodiversity on Earth.

All Nordic Journal of Botany volumes since the journal started in 1981 are available as fulltext  
for subscribers online at Wiley Online Library – [click here](#)

To take out a personal subscription, please [click here](#)

More information about Nordic Journal of Botany at <http://nordicjbotany.org>

## *Gymnosporia thyrsoflora* comb. nov. (Celastraceae), a correct name to replace *G. graciliramula* from southwest China

Wen-Bin Yu, Hong Wang and De-Zhu Li

W.-B. Yu (yuwenbin@mail.kib.ac.cn), H. Wang, D.-Z. Li, Key Laboratory of Biodiversity and Biogeography, Kunming Inst. of Botany, Chinese Academy of Sciences, 132 Lanhei Road, Kunming, CN-650204 Yunnan, China. WBY also at Dept of Botany, Field Museum of Natural History, 1400 S Lake Shore Drive, Chicago, US-60605 Illinois, USA.

In 'Flora of China', the combination '*Gymnosporia graciliramula* (S. J. Pei & Y. H. Li) Q. R. Liu & Funston' was proposed, but this name was not validly published in 2008 because the presumed basionym *Maytenus graciliramula* S. J. Pei & Y. H. Li was not published in 1979 as stated, but in 1981, contrary to Article 33.4 of ICBN. Meanwhile, two older names, *M. thyrsoflora* S. J. Pei & Y. H. Li and *M. pseudoracemosa* S. J. Pei & Y. H. Li, were listed as synonyms in conflict with article 52.3 of ICBN. In this article, we propose a new combination *G. thyrsoflora* (S. J. Pei & Y. H. Li) W. B. Yu & D. Z. Li that is the correct name when *G. thyrsoflora*, *M. pseudoracemosa* and *M. graciliramula* are treated as a single species in *Gymnosporia*.

The genus *Gymnosporia* was first established by Hooker (1862) on the basis of *Celastrus* sect. *Gymnosporia* Wight and Arnott. In the last century, nevertheless, several taxonomists suggested that *Gymnosporia* should be congeneric with *Maytenus* Molina s.l. (Pei and Li 1981a, Jordaan and van Wyk 1999). However, recently the genus *Gymnosporia* was reinstated as distinct from *Maytenus* s.s. based on comparative analyses of morphological characters and molecular sequences (Jordaan and van Wyk 1999, Simmons and Hedin 1999, Simmons et al. 2001a, 2001b, 2008). Meanwhile, McKenna et al. (2011) stated that the current circumscription of *Gymnosporia* included the Old World *Gymnosporia* and *Putterlickia* Endl., and the New World species of *Gymnosporia* were transferred to a new genus *Haydenia* M. P. Simmons.

Previously, Chinese taxonomists treated *Gymnosporia* as included in *Maytenus* s.l. (Pei and Li 1981a, Cheng et al. 1999, Qin et al. 2008). During the revision of *Maytenus* s.l. from Yunnan, Pei and Li (1979) described ten new taxa, including *M. graciliramula*, *M. thyrsoflora* and *M. pseudoracemosa*. However, the name *M. graciliramula* was not validly published because of two collections simultaneously designated as type (Article 40.1 and 40.2). *Maytenus graciliramula* became validated later (Pei and Li 1981b). In the Chinese edition of 'Flora Reipublicae Popularis Sinicae', Cheng et al. (1999) adopted *Maytenus* sect. *Gymnosporia* (Wight & Arnott) C. Y. Cheng, and they accepted *M. thyrsoflora* and treated *M. pseudoracemosa* and *M. graciliramula* as its synonyms. According to Article 11.5, this synonymization

established priority of the name *M. thyrsoflora* over the name *M. pseudoracemosa*. During the preparation of the updated and English edition of 'Flora of China', Liu and Funston (2008) recognized *Gymnosporia* as a distinct genus with three new combinations proposed. Unfortunately, their intended combination '*G. graciliramula* (S. J. Pei & Y. H. Li) Q. R. Liu & Funston' was not validly published, because they only provided a full and direct reference to the publication in 1979 contrary to Article 41.1 of ICBN (McNeill et al. 2012). Meanwhile, Liu and Funston (2008) listed two older names, *M. thyrsoflora* and *M. pseudoracemosa*, as synonyms of their '*G. graciliramula*' but without full and direct references to the places of their valid publication, therefore not making the intended new combination validly published even as a superfluous replacement name. The new combination *G. thyrsoflora* (S. J. Pei & Y. H. Li) W. B. Yu & D. Z. Li proposed herein becomes the correct name when *G. thyrsoflora*, *M. pseudoracemosa* and *M. graciliramula* are united in *Gymnosporia*.

***Gymnosporia thyrsoflora* (S. J. Pei & Y. H. Li) W. B. Yu & D. Z. Li comb. nov.**

**Basionym:** *Maytenus thyrsoflora* S. J. Pei & Y. H. Li in Res. Bull. Trop. Plants 13 (1979, p. 13).

**Type:** China. Yunnan: Shuangjiang, 24 Jul 1977, J. H. Zhang and Y. Cheng 11054 (holotype: HITBC 081516!, isotypes: HITBC 081517!, 081518!).

**Taxonomic synonyms:** *Maytenus graciliramula* S. J. Pei & Y. H. Li (1981, pp. 242–243); ‘*Maytenus graciliramula* S. J. Pei & Y. H. Li’ (1979, p. 13), not validly published (Article 40.1 and 40.2); ‘*Gymnosporia graciliramula* (S. J. Pei & Y. H. Li) Q. R. Liu and Funston’ (2008, p. 476), not validly published (Article 41.1).

**Type:** China. Yunnan: Linchang, 19 Jul 1977, J. H. Zhang and Y. Cheng 11050 (holotype: HITBC 081504!).

– *Maytenus pseudoracemosa* S. J. Pei & Y. H. Li (1979, p. 12).

**Type:** China. Yunnan: Mengla, 23 May 1976, G. D. Tao 9113 (holotype: HITBC 081511!, isotype: HITBC 081512!).

*Gymnosporia thyrsoflora* is endemic to southwest China (southern Yunnan and western Guangxi). It occurs on open dry slopes with mixed shrubs along riversides at altitudes between 400 and 1500 m a.s.l. Morphologically, *G. thyrsoflora* belongs to *G.* sect. *Tenuispinae* in accordance with Jordaan and van Wyk (2006). This species is distinguished from the other Chinese *Gymnosporia* in having axillary and slender thorns on young branches and sturdy thorns on old branches, narrowly obovate and papery leaves, 3-valved red–brown capsules, and a white cupulate aril covering the base of seeds.

*Acknowledgements* – We are grateful to the curator of the herbarium of HITBC for making specimens available; and to Dr Gerry Moore for valuable comments and suggestions. This study was supported by the West Light Foundation of Chinese Academy of Sciences.

## References

Cheng, C.-Y. et al. 1999. Celastraceae (excluding *Dipentodon* and *Perrottetia*). – In: Cheng, C.-Y. and Huang, P.-H. (eds), *Flora Reipublicae Popularis Sinicae*. Science Press, pp. 1–218.

Hooker, J. D. 1862. Celastrineae. – In: Bentham, G. and Hooker, J. D. (eds), *Genera plantarum*. Vol. 1. Reeve, pp. 357–371.

Jordaan, M. and van Wyk, A. E. 1999. Systematic studies in subfamily Celastroideae (Celastraceae) in southern Africa: reinstatement of the genus *Gymnosporia*. – *S. Afr. J. Bot.* 65: 177–181.

Jordaan, M. and van Wyk, A. E. 2006. Sectional classification of *Gymnosporia* (Celastraceae), with notes on the nomenclatural and taxonomic history of the genus. – *Taxon* 55: 515–525.

Liu, Q.-R. and Funston, A. M. 2008. *Gymnosporia* (Wight & Arnott) Bentham & J. D. Hooker. – In: Wu, Z.-Y. et al. (eds), *Flora of China*. Science Press; Miss. Bot. Gard. Press, pp. 474–477.

McKenna, M. J. et al. 2011. Delimitation of the segregate genera of *Maytenus* s. l. (Celastraceae) based on morphological and molecular characters. – *Syst. Bot.* 36: 922–932.

McNeill, J. et al. (eds) 2012. International Code of Nomenclature for algae, fungi, and plants (Melbourne Code) adopted by the Eighteenth International Botanical Congress Melbourne, Australia, July 2011. *Regnum Vegetabile* 154: I–XXX + 1–208.

Pei, S.-J. and Li, Y.-H. 1979. The genus *Maytenus* Molina From Yunnan. – *Res. Bull. Trop. Plants* 13: 11–22.

Pei, S.-J. and Li, Y.-H. 1981a. A taxonomical problem of the genus *Maytenus* Molina and *Gymnosporia* (Wight & Arn.) Benth & Hook. f. from China. – *Acta Bot. Yunnan.* 3: 25–31.

Pei, S.-J. and Li, Y.-H. 1981b. New materials for Yunnan *Maytenus* Molina. – *Acta Bot. Yunnan.* 3: 239–248.

Qin, X.-S. et al. 2008. A new species of *Maytenus* section *Gymnosporia* (Celastraceae) from Hainan Island, China. – *Bot. J. Linn. Soc.* 158: 534–538.

Simmons, M. P. and Hedin, J. P. 1999. Relationships and morphological character change among genera of Celastraceae sensu lato (including Hippocrateaceae). – *Ann. Miss. Bot. Gard.* 86: 723–757.

Simmons, M. P. et al. 2001a. Phylogeny of the Celastraceae inferred from phytochrome B gene sequence and morphology. – *Am. J. Bot.* 88: 313–325.

Simmons, M. P. et al. 2001b. Phylogeny of the Celastraceae inferred from 26S nuclear ribosomal DNA, phytochrome B, *rbcL*, *atpB*, and morphology. – *Mol. Phylogen. Evol.* 19: 353–366.

Simmons, M. P. et al. 2008. Phylogeny of the Celastreae (Celastraceae) and the relationships of *Catha edulis* (qat) inferred from morphological characters and nuclear and plastid genes. – *Mol. Phylogen. Evol.* 48: 745–757.